

# NONNEGATIVE INTERPOLATION\*

SHI YING-GUANG (史应光)

(Computing Center, Academia Sinica, Beijing, China)

## Abstract

The problem discussed in this paper is to determine a nonnegative interpolating polynomial which takes the prescribed nonnegative values  $y_0, y_1, \dots, y_n$  at given distinct points  $x_0, x_1, \dots, x_n$ :

$$p(x_i) = y_i, \quad i=0, 1, \dots, n.$$

This paper shows: (1)  $2n$  is the least number of  $m$  such that there exists a polynomial  $p \in P_m^+$ , the set of all nonnegative polynomials of degree  $\leq m$ , satisfying the above equations for any choice of  $y_i \geq 0$ . (2) The above equations have a unique solution in  $P_{2n}^+$  if and only if at most one of the  $y_i$ 's is nonzero.

## 1. Introduction

Let  $x_0, x_1, \dots, x_n$  be  $n+1$  distinct points in the interval  $[a, b]$  satisfying

$$a \leq x_0 < x_1 < \dots < x_n \leq b.$$

Let  $y_0, y_1, \dots, y_n$  be any prescribed values satisfying  $y_i \geq 0, i=0, 1, \dots, n$ . Let  $P_m$  denote the set of all polynomials of degree equal to  $m$  or less. In this paper we consider the following problem: to find a nonnegative polynomial  $p$  such that

$$p(x_i) = y_i, \quad i=0, 1, \dots, n. \quad (1)$$

That is to say, if we denote by  $P_m^+$  the set of all nonnegative polynomials of degree equal to  $m$  or less, then our problem is to determine a polynomial  $p \in P_m^+$  satisfying the above interpolating conditions.

It is not hard to see that the problem is not necessarily solvable for  $m=n$ . For example, taking  $n=1, x_0=a, x_1=\frac{1}{2}(a+b), y_0=1$  and  $y_1=0$ , there does not exist a  $p \in P_1^+$  satisfying (1). Therefore we are concerned with the existence of a nonnegative polynomial satisfying (1), in particular, the least number of  $m$  for which the problem is always solvable for any choice of  $y_i \geq 0, i=0, 1, \dots, n$ . That is the content of Section 2. Section 3 will discuss uniqueness of such a polynomial.

## 2. Existence

The main result in this section is as follows.

**Theorem 1.**  $2n$  is the least number of  $m$  such that there exists a polynomial  $p \in P_m^+$  satisfying the equations (1) for any choice of  $y_i \geq 0, i=0, 1, \dots, n$ .

*Proof.* First, we are going to show that there exists a polynomial  $p \in P_{2n}^+$  satisfying (1) for any choice of  $y_i \geq 0$ . To do this, put

$$w(x) = (x-x_0)(x-x_1)\dots(x-x_n),$$

\* Received December 5, 1983.

$$l_i(x) = \frac{w(x)}{(x-x_i)w'(x_i)}, \quad i=0, 1, \dots, n,$$

here

$$w'(x_i) = \prod_{\substack{j=1 \\ j \neq i}}^n (x_i - x_j), \quad i=0, 1, \dots, n.$$

Then

$$p(x) = \sum_{i=0}^n y_i l_i^2(x) \tag{2}$$

is such a polynomial.

In fact, it is clear that  $p \in P_{2n}^+$ . On the other hand, by an observation one can see that

$$l_i(x_j) = \begin{cases} 1, & j=i, \\ 0, & j \neq i, \end{cases}$$

Hence

$$p(x_i) = y_i, \quad i=0, 1, \dots, n.$$

We turn now to show the minimality of  $m=2n$ . To the end let us consider some choice of  $y_i$ :

$$y_0=1, \quad y_1=\dots=y_n=0.$$

Suppose that  $p \in P_m^+$  satisfies

$$p(x_i) = y_i, \quad i=0, 1, \dots, n,$$

i. e.,

$$p(x_0)=1, \quad p(x_1)=\dots=p(x_n)=0.$$

Since  $p \geq 0$ , each of  $x_1, \dots, x_n$  is a zero of at least multiplicity 2 of  $p$ . Meanwhile  $p \neq 0$ , then  $p$  should contain a factor  $(x-x_1)^2 \dots (x-x_n)^2$ . This means that  $p$  has degree at least  $2n$  and therefore  $m \geq 2n$ . This proves the minimality of  $m=2n$ .

### 3. Uniqueness

A simple example indicates that a solution of the interpolatory problem in  $P_{2n}^+$  is not unique for some choice of  $y_i \geq 0, i=0, 1, \dots, n$ .

*Example.* Let  $n=2, [a, b] \equiv [-1, 1], x_0=-1, x_1=0, x_2=1, y_0=1, y_1=0,$  and  $y_2=1$ . Both  $x^2$  and  $x^4$  are the polynomials in  $P_4^+$  satisfying (1). Generally,  $p_t = tx^4 + (1-t)x^2$  with  $0 \leq t \leq 1$  is also such a polynomial.

Precisely, we have the following criterion of uniqueness of a solution of the interpolatory problem in  $P_{2n}^+$ .

**Theorem 2.** *The polynomial (2) is the unique one satisfying the equations (1) from  $P_{2n}^+$  if and only if at most one of the  $y_i$ 's is nonzero.*

*Proof.* Sufficiency. Suppose that, say,  $y_i=0, \forall i \neq k$  for some index  $k$ . Thus

$$p(x) = y_k l_k^2(x).$$

Let  $q \in P_{2n}^+$  satisfy

$$q(x_i) = y_i, \quad i=0, 1, \dots, n$$

for such a choice of the  $y_i$ 's. Then we obtain

$$p(x_i) = q(x_i), \quad i=0, 1, \dots, n$$

and  $p'(x_i) = q'(x_i) = 0, \quad i=0, \dots, k-1, k+1, \dots, n.$

Noting that  $p, q \in P_{2n}$ , this yields that  $p=q$ .

*Necessity.* Denote  $I = \{i: y_i > 0\}$  and  $J = \{j: y_j = 0\}$ . Suppose, on the contrary, that  $\text{card } I > 1$ , where  $\text{card } I$  denotes the cardinality of  $I$ . Write  $s = \text{card } J$ . Now let  $q \in P_{2n}$  satisfy