

Enhancing Jerk Nonlinear Equation by Adomain Decomposition Method

Abdulrahmn N. Akour^{1,†}, Emad K. Jaradat², Hadeel Saqa-Allah³ and
Omar K. Jaradat⁴

Abstract In this manuscript, an approximate periodic solution of a nonlinear Jerk equation concerning the third order time derivative (ODE) is demonstrated. This method succeeds highly in constructing a periodic solution for Jerk nonlinear differential equations. The results investigated from this method are compared with those determined from other analytical and numerical methods. Our results demonstrate a significant consistency with the numerical solution in addition to the analytical methods.

Keywords Adomian decomposition method, nonlinear Jerk equation, high order differential equation

MSC(2010) 34A34 , 70Kxx.

1. Introduction

Jerk equation (JE) expresses the acceleration change with time; JE is the first derivative of acceleration (j) and can be expressed as follows.

$$\vec{j} = \frac{d\vec{a}}{dt} = \frac{d^2\vec{v}}{dt^2} = \frac{d^3\vec{s}}{dt^3}. \quad (1.1)$$

Having that

$\vec{a}$ is acceleration;

$\vec{v}$ is velocity;

$\vec{s}$ is displacement;

t is time.

As JE is a third derivative of displacement, differential equation (DE) of the form

$$J(\ddot{x}, \ddot{x}, \dot{x}, x) = 0 \quad (1.2)$$

Known as JE. It was found that a JE, which describes a system formed of three first-order, ordinary non-linear DE, is the minimal expression showing a chaotic

[†]the corresponding author.

Email address:abd-akour@bau.edu.jo, abdulrahmanakour1@yahoo.com. (A. Akour), ekyjaradat@imamu.edu.sa.(E. Jaradat), hadeelsaqallah@gmail.com. (H. Saqa-Allah), jaradat@mutah.edu.jo. (O. Jaradat)

¹Department of Basic Scientific Sciences, Al-Huson College, Al-Balqa Applied University, Salt, Jordan

²Department of Physics, Faculty of Science, Imam Mohammad Ibn Saud Islamic University, Riyadh, Saudi Arabia

³Department of physics, Mutah University, Al-Karak, Jordan

⁴Department of mathematics, Mutah University, Al-Karak, Jordan

manner [1]. The familiar form of the JE that keeps its style upon time and space reversal, and with the cubic non-linearities is determined as:

$$\ddot{x} = -\gamma\dot{x} - \alpha\dot{x}^3 - \beta x^2\dot{x} + \delta x\dot{x}\ddot{x} - \epsilon\dot{x}\ddot{x}^2, \quad (1.3)$$

where the parameters ϵ , δ , β , α and γ are constant. Here, for special cases and to compare with other studies at least one of the parameters δ , β and ϵ should be not equal to zero. The initial conditions (IC) are [1]

$$x(0) = 0, \quad \dot{x}(0) = B \neq 0, \quad \ddot{x}(0) = 0. \quad (1.4)$$

JE spreads widely in several mechanics and acoustics applications. Therefore, many methods are demonstrated to investigate an exact or approximate solution for it. Modified harmonic balance method (MHBM) [2] is used to find a 2^{nd} approximate solution of non-linear, third-order differential JE, but it is successful for a limited number of other non-linear equations. Hybrid block method (HBM) [3] solves JE with one hybrid point that is indicated to enhance effectively the local truncation errors of the basic formulas of the solutions and the derivatives at the end of the block. The Homotopy method, considered as one of the powerful methods in many physical and mechanical approximations [4], is extended to provide periodic solutions of a nonlinear jerk equation involving the 3^{rd} order time derivative. A suitable parameter is introduced to provide a numerical solution that proves well for the first few terms. The Analytical solutions [5], by Ramose, are developed for nonlinear, 3^{rd} order nonlinear DE to investigate a convergent solution. The solutions are determined as functions of the coefficients that concern the terms that are linearly proportional to the velocity and nonlinear terms, using means of transformations for periodic and non-periodic behavior. In our study, we use the LDM. The LDM is an effective, and accurate method for demonstrating analytical solutions to non-linear problems. Also, it competes highly with other methods in finding accurate or approximate solutions for different nonlinear equations of physical or mechanical applications [6]. On the other hand, LDM proves the highest advantage in exceeding the difficulty that appears in the calculation of the Adomain polynomials, compared with other methods. The LDM divides the given equation into non-linear and linear parts, next obtains the highest-order derivative operator incorporated with the linear operator on each side, then provides Adomian's polynomials, and eventually investigates the series successive terms of the solution by a frequent relation using Adomian's polynomials. [7].

$$Lu + Nu + Ru = g \quad (1.5)$$

where L is an operator representing the highest order derivative; R is a linear operator for the remainder of the linear portion; N(u) represents the non-linear terms; lastly, g is the source term. This article is divided into three sections: the current section which demonstrates the introduction; the second section which demonstrates the mathematical analysis and results; and lastly, the conclusion in section 3.

2. Mathematical analysis and results

With appreciation of the general case, let us consider the case for $\alpha = \beta = 1$ and $\gamma = \delta = \epsilon = 0$. So, Eq.(1.3) becomes

$$\ddot{x} = -\dot{x}^3 - x^2\dot{x}. \quad (2.1)$$