

Bifurcations and Exact Solutions of the Gerdjikov-Ivanov Equation*

Jinsen Zhuang¹, Yan Zhou^{1,†} and Jibin Li^{1,2}

Abstract For the Gerdjikov-Ivanov equation, by using the method of dynamical system, this paper investigates the exact explicit solutions with the form $q(x, t) = \phi(\xi) \exp[i(\kappa x - \omega t + \theta(\xi))]$, $\xi = x - ct$. In the given parameter regions, more than 14 explicit exact parametric representations are presented.

Keywords Bifurcation, exact solution, planar Hamiltonian system, Gerdjikov-Ivanov equation

MSC(2010) 34C23, 35Q51, 35Q52, 35Q53, 58J55.

1. Introduction

In [8], Fang stated that “The nonlinear Schrödinger (NLS) equation is one of the most generic soliton equations, and arises from a wide variety of fields such as quantum field theory, weakly nonlinear dispersive water waves and nonlinear optics. To study the effect of higher-order perturbations, various modifications and generalizations of the NLS equations have been proposed and studied. Among them, there are three celebrated equations with derivative-type nonlinearities, which are called the derivative nonlinear Schrödinger (DNLS) equations. One is the Kaup-Newell equation [17]:

$$iq_t + q_{xx} + i(|q|^2 q)_x = 0, \quad (1.1)$$

which is usually called DNLSI. The second type is the Chen-Lee-Liu equation [4]:

$$iq_t + q_{xx} + i|q|^2 q_x = 0, \quad (1.2)$$

which is called DNLSII. The last one takes the form [13]:

$$iq_t + q_{xx} - iq^2 q_x^* + \frac{1}{2}q^3 (q^*)^2 = 0, \quad (1.3)$$

which is called the Gerjikov–Ivanov (GI) equation or DNLSIII. In equation (1.3), q^* denotes the complex conjugation of q .

[†]The corresponding author.

Email address: zzjinsen@hqu.edu.cn (J. Zhuang), zy4233@hqu.edu.cn (Y. Zhou), lijb@zjnu.cn (J. Li)

¹School of Mathematical Sciences, Huaqiao University, Quanzhou, Fujian 362021, China

²Department of Mathematics, Zhejiang Normal University, Jinhua, Zhejiang 321004, China

*The authors were supported by National Natural Science Foundations of China (Nos. 11871231, 12071162, 11701191) and the Natural Science Foundation of Fujian Province (No. 2021J01303).

Equation (1.3) has been studied by many authors (see [1, 2, 5–7, 9–12, 14–16, 18, 21–32]). The purpose of this paper is to study some new exact traveling wave solutions in explicit form of the GI equation by using the bifurcation theory of dynamical system. We assume that the exact solutions of equation (1.3) take the form:

$$q(x, t) = \phi(\xi) \exp [i(\kappa x - \omega t + \theta(\xi))], \quad \xi = x - ct, \quad (1.4)$$

where c is the wave velocity, and $\phi(\xi), \theta(\xi)$ are two functions with variable ξ , κ and ω are two constant parameters. Substituting (1.4) into equation (1.3) and separating the real and imaginary parts respectively, we have

$$\phi'' + (c - 2\kappa)\theta'\phi - (\kappa + \theta')\phi^3 + (\omega - \kappa^2)\phi - (\theta')^2\phi = 0, \quad (1.5)$$

$$\phi'\phi^2 - \theta''\phi - 2\phi'\theta' + (c - 2\kappa)\phi' = 0, \quad (1.6)$$

where “'” is the derivative with respect to ξ . Integrating (1.6), it follows that $(c - 2\kappa)\phi + \frac{1}{3}\phi^3 = C_1 + \theta'\phi + \int \theta' d\phi$, where C_1 is an integral constant. Thus, we obtain $C_1 = 0$ and

$$\theta' = \frac{1}{2}(c - 2\kappa) + \frac{1}{4}\phi^2, \quad \theta(\xi) = \frac{1}{2}(c - 2\kappa)\xi + \frac{1}{4} \int \phi^2(\xi) d\xi. \quad (1.7)$$

Substituting (1.7) into (1.5), we obtain the following planar dynamical system:

$$\frac{d\phi}{d\xi} = y, \quad \frac{dy}{d\xi} = - \left(\left(\frac{1}{4}c^2 - c\kappa + \omega \right) \phi - \frac{1}{2}c\phi^3 + \frac{3}{16}\phi^5 \right). \quad (1.8)$$

System (1.8) has the first integral:

$$H(\phi, y) = \frac{1}{2}y^2 + \frac{1}{2}\alpha\phi^2 - \frac{c}{8}\phi^4 + \frac{1}{32}\phi^6 = h, \quad (1.9)$$

where $\alpha = \frac{1}{4}c^2 - c\kappa + \omega$.

System (1.8) is a three-parameter planar dynamical system depending on the parameter group (c, κ, ω) . Since the parametric representations of the phase orbits defined by the vector fields of system (1.8) give rise to all exact solutions with the form (1.4) of equation (1.3), we need to investigate the bifurcations of phase portraits for system (1.8) in the (ϕ, y) -phase plane as the parameters are changed (see [19, 20, 33]).

The main result in the present paper is summarized as follows.

Theorem 1.1. *Assume that the parameter $c > 0$ of system (1.8) is fixed. Consider the solutions of equation (1.3) with the form $q(x, t) = \phi(\xi) \exp [i(\kappa x - \omega t + \theta(\xi))]$. Then, the following conclusions hold.*

(i) *For any $(\kappa, \omega) \in R^2$ in Figure 1 (a), corresponding to the families of the periodic orbits of system (1.8), equation (1.1) always has the exact explicit solutions with the parametric representations given by (4.1) or (4.2).*

(ii) *When $(\kappa, \omega) \in (II), (III), (IV)$ and $(L_2), (L_3)$ in Figure 1 (a), corresponding to the families of the periodic orbits of system (1.8), equation (1.1) has the exact explicit solutions with the parametric representations given by (4.3), (4.4), (4.5), (4.6) and (4.7).*

(iii) *When $(\kappa, \omega) \in (II), (III)$ and $(L_1), (L_2)$ in Figure 1 (a), corresponding to the heteroclinic orbits of system (1.8), equation (1.1) has the exact explicit solutions with the parametric representations given by (4.8), (4.9) and (4.10).*

(iv) When $(\kappa, \omega) \in (II), (III), (IV)$ and $(L_2), (L_3)$ in Figure 1 (a), corresponding to the homoclinic orbits of system (1.8), equation (1.1) has the exact explicit solutions with the parametric representations given by (4.11), (4.12), (4.13) and (4.14).

The proof of this theorem is given in the following Sections 2, 3 and 4.

The rest of this paper is organized as follows: in Section 2, the bifurcations of the phase portraits of system (1.8) are studied. In Section 3, all exact explicit parameter representations of bounded phase orbits of system (1.8) are given in given parameter regions. In Section 4, corresponding to the solutions of system (1.8), all exact explicit parameter representations of equation (1.3) are derived.

2. Bifurcations of phase portraits of system (1.8)

To find the equilibrium points of system (1.8), write that $f(\phi) = \alpha - \frac{1}{2}c\phi^2 + \frac{3}{16}\phi^4$. Clearly, if $\Delta = c^2 + 12(c\kappa - \omega) > 0$, then when $\phi^2 = \frac{2}{3}(2c \mp \sqrt{\Delta})$, $f(\phi) = 0$. Thus, for a fixed $c > 0$, we have the following conclusions.

- (i) When $\omega > c\kappa + \frac{1}{12}c^2$, $\Delta < 0$, system (1.8) has only one singular point $O(0, 0)$.
- (ii) When $\omega = c\kappa + \frac{1}{12}c^2$, $\Delta = 0$, system (1.8) has one simple singular point $O(0, 0)$ and two double equilibrium points $E_{1,2}(\mp\frac{2}{3}\sqrt{3c}, 0)$.
- (iii) When $c\kappa - \frac{1}{4}c^2 < \omega < c\kappa + \frac{1}{12}c^2$, we have $\Delta > 0$, $2c - \sqrt{\Delta} > 0$. Hence, on the ϕ -axis, there exist five equilibrium points of system (1.8) at $E_1(-\phi_2, 0)$, $E_2(-\phi_1, 0)$, $O(0, 0)$, $E_3(\phi_1, 0)$ and $E_4(\phi_2, 0)$, where $\phi_{1,2} = \left(\frac{2}{3}(2c \mp \sqrt{\Delta})\right)^{\frac{1}{2}}$.
- (iv) When $\omega = c\kappa - \frac{1}{4}c^2$, we have $2c - \sqrt{\Delta} = 0$. system (1.8) has a high-order equilibrium point $O(0, 0)$ and two simple equilibrium points $E_1(-\phi_2, 0)$ and $E_2(\phi_2, 0)$.
- (v) When $\omega < c\kappa - \frac{1}{4}c^2$, system (1.8) has three simple equilibrium points at $O(0, 0)$, $E_1(-\phi_2, 0)$ and $E_2(\phi_2, 0)$.

Let $M(\phi_j, 0)$ be the coefficient matrix of the linearized system of system (1.8) at the equilibrium point E_j . Let $J(\phi_j, 0)$ be its Jacobin determinant. Then, one has

$$J(0, 0) = \alpha, \quad J(\phi_j, 0) = \phi_j f'(\phi_j).$$

By the theory of planar dynamical system, for an equilibrium point of a planar Hamiltonian system, if $J < 0$, then the equilibrium point is a saddle point; if $J > 0$, then it is a center point; if $J = 0$ and the Poincaré index of the equilibrium point is 0, then this equilibrium point is a cusp.

Now, write that $h_j = H(\phi_j, 0)$, $h_0 = H(0, 0) = 0$. We have that $h_1 = \frac{\Delta}{54}(c + \sqrt{\Delta})$, $h_2 = \frac{\Delta}{54}(c - \sqrt{\Delta})$. Obviously, when $\omega = c\kappa$, we have $h_2 = 0$.

Based on the above results, for a fixed parameter $c > 0$, in the (κ, ω) -parameter plane, there exist three bifurcation straight lines $(L_1) : \omega = c\kappa + \frac{1}{12}c^2$; $(L_2) : \omega = c\kappa$; and $(L_3) : \omega = c\kappa - \frac{1}{4}c^2$. These straight lines divide the (κ, ω) -parameter plane into the four regions: (I), (II), (III), (IV) (see Figure 1 (a)). As ω is varied, we obtain the bifurcations of the phase portraits of system (1.8), as shown in Figure 1(b)-(h).




Figure 1. Bifurcations of phase portraits of system (1.8) for a fixed $c > 0$

3. Exact parametric representations of the solutions defined by all bounded orbits of system (1.8)

We see from (1.9) that

$$y^2 = 2h - \alpha\phi^2 + \frac{c}{4}\phi^4 - \frac{1}{16}\phi^6. \tag{3.1}$$

By using the first equation of (1.8), we obtain

$$\xi = \int_{\phi_0}^{\phi} \frac{4d\phi}{\sqrt{16h - 8\alpha\phi^2 + 4c\phi^4 - \phi^6}} = \int_{\psi_0}^{\psi} \frac{2d\psi}{\sqrt{\psi(16h - 8\alpha\psi + 4c\psi^2 - \psi^3)}}. \tag{3.2}$$

Thus, we can calculate the exact parametric representations defined by the bounded orbits of system (1.8).

3.1. Exact periodic solutions of system (1.8) when $(\kappa, \omega) \in (I)$ in Figure 1(a)

In this case, corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, \infty)$, there exists a family of periodic orbits of system (1.8) (see Figure 1(b)). Now, (3.2) can be written as $\frac{1}{2}\xi = \int_0^{\psi} \frac{d\psi}{\sqrt{(\psi_1 - \psi)\psi(\psi + \psi_3)(\psi + \psi_4)}}$ or $\frac{1}{2}\xi = \int_0^{\psi} \frac{d\psi}{\sqrt{(\psi_1 - \psi)\psi[(\psi - b_1)^2 + a_1^2]}}$. Hence, we have the following two exact parametric representations of the periodic family of system (1.8):

$$\phi(\xi) = \frac{\sqrt{\psi_3}\hat{\alpha}_1 \operatorname{sn}(\Omega_1 \xi, k)}{(1 - \hat{\alpha}_1^2 \operatorname{sn}^2(\Omega_1 \xi, k))^{\frac{1}{2}}}, \tag{3.3}$$

where $\hat{\alpha}_1^2 = \frac{\psi_1}{\psi_1 + \psi_3}$, $k^2 = \frac{\hat{\alpha}_1^2(\psi_4 - \psi_3)}{\psi_4}$, $\Omega_1 = \frac{1}{4}\sqrt{(\psi_1 + \psi_3)\psi_4}$, $\text{sn}(u, k)$, $\text{cn}(u, k)$, $\text{dn}(u, k)$ are the Jacobian elliptic functions (see Byrd and Fridman [3]).

$$\phi(\xi) = \left(\alpha_1 + \frac{\beta_1}{1 + \hat{\alpha}_2 \text{cn}(\Omega_2 \xi, k)} \right)^{\frac{1}{2}}, \tag{3.4}$$

where $A_1^2 = (\psi_1 - b_1)^2 + a_1^2$, $B_1^2 = b_1^2 + a_1^2$, $k^2 = \frac{(\psi_1^2 - (A_1 - B_1)^2)}{4A_1 B_1}$, $\alpha_1 = \frac{-\psi_1 B_1}{A_1 - B_1}$, $\beta_1 = \frac{2A_1 B_1}{A_1^2 - B_1^2}$, $\hat{\alpha}_2 = \frac{A_1 - B_1}{A_1 + B_1}$, $\Omega_2 = \frac{1}{2}\sqrt{A_1 B_1}$.

3.2. Exact periodic solutions of system (1.8) when $(\kappa, \omega) \in (L_1)$ in Figure 1(a).

(i) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, h_1), (h_1, \infty)$, $h_1 = h_2 = \frac{2c^3}{27}$, there exist two families of periodic orbits of system (1.8) (see Figure 1(c)). These periodic orbits have the same exact parametric representations as (3.4).

(ii) Corresponding to the level curves defined by $H(\phi, y) = \frac{2c^3}{27}$, there exist two heteroclinic orbits connecting two double equilibrium points $E_{1,2}(\mp \frac{2}{3}\sqrt{3c}, 0)$ of system (1.8) (see Figure 1(c)). Now, (3.2) can be written as $\xi = \int_0^\phi \frac{12\sqrt{3}d\phi}{(4c - 3\phi^2)^{\frac{3}{2}}} = \frac{3\sqrt{3}\phi}{c\sqrt{4c - 3\phi^2}}$. Thus, it follows the exact parametric representations:

$$\phi(\xi) = \mp \frac{4c^{\frac{3}{2}}\xi}{\sqrt{3(9 + c^2\xi^2)}}. \tag{3.5}$$

3.3. Exact periodic solutions and homoclinic, heteroclinic solutions of system (1.8) when $(\kappa, \omega) \in (II)$ in Figure 1(a).

In this case, we have $0 < h_2 < h_1$.

(i) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, h_2]$, there exists a family of periodic orbits of system (1.8), enclosing the origin $O(0, 0)$ (see Figure 1(d)). When $0 < h < h_2$, these periodic orbits have the same exact parametric representations as (3.4).

(ii) Corresponding to the level curves defined by $H(\phi, y) = h, h \in [h_2, h_1)$, there exist three families of periodic orbits of system (1.8), enclosing the origin $O(0, 0)$, $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively (see Figure 1(d)). Now, (3.2) can be written as $\frac{1}{2}\xi = \int_0^\psi \frac{d\psi}{\sqrt{(\psi_a - \psi)(\psi_b - \psi)(\psi_c - \psi)\psi}}$ and $\frac{1}{2}\xi = \int_{\psi_b}^\psi \frac{d\psi}{\sqrt{(\psi_a - \psi)(\psi - \psi_b)(\psi - \psi_c)\psi}}$. These periodic orbits have the exact parametric representations as follows:

$$\phi(\xi) = \frac{\sqrt{\psi_a}|\hat{\alpha}_3|\text{sn}(\Omega_3 \xi, k)}{(1 - \hat{\alpha}_3^2 \text{sn}^2(\Omega_3 \xi, k))^{\frac{1}{2}}}, \tag{3.6}$$

where $\hat{\alpha}_3^2 = \frac{-\psi_c}{\psi_a - \psi_c}$, $k^2 = \frac{-\hat{\alpha}_3^2(\psi_a - \psi_b)}{\psi_b}$, $\Omega_3 = \frac{1}{4}\sqrt{(\psi_a - \psi_c)\psi_b}$ and

$$\phi(\xi) = \mp \left(\psi_c + \frac{\psi_b - \psi_c}{1 - \hat{\alpha}_4^2 \text{sn}^2(\Omega_3 \xi, k)} \right)^{\frac{1}{2}}, \tag{3.7}$$

where $\hat{\alpha}_4^2 = \frac{\psi_a - \psi_b}{\psi_a - \psi_c}$, $k^2 = \frac{\hat{\alpha}_4^2 \psi_c}{\psi_b}$.

Notice that the level curve defined by $H(\phi, y) = h_2$ contain a periodic orbit enclosing the origin $O(0, 0)$ and two singular points $E_1(-\phi_2, 0), E_4(\phi_2, 0)$. In this case, in (3.6), $\psi_a = \psi_b = \phi_2^2, k^2 = 0, \hat{\alpha}_3^2 = \frac{-\psi_c}{\phi_2^2 - \psi_c}$ and $\Omega_3 = \frac{1}{4}\sqrt{(\phi_2^2 - \psi_c)\phi_2^2}$. The periodic orbit has the exact parametric representation:

$$\phi(\xi) = \frac{\phi_2|\hat{\alpha}_3|\sin(\Omega_3\xi)}{(1 - \hat{\alpha}_3^2 \sin^2(\Omega_3\xi))^{\frac{1}{2}}}. \tag{3.8}$$

(iii) Corresponding to the level curves defined by $H(\phi, y) = h_1$, there exist two homoclinic orbits enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively, and two heteroclinic orbits enclosing the origin $O(0, 0)$ (see Figure 1 (d)). In this case, (3.2) becomes that $\frac{1}{2}\xi = \int_0^\psi \frac{d\psi}{(\psi_1 - \psi)\sqrt{(\psi_M - \psi)\psi}}$ and $\frac{1}{2}\xi = \int^\psi \frac{d\psi}{(\psi - \psi_1)\sqrt{(\psi_M - \psi)\psi}}$, where $\psi_1 = \phi_1^2 = \frac{2}{3}(2c - \sqrt{\Delta}), \psi_M = \frac{4}{3}(c + \sqrt{\Delta})$. Thus, the two heteroclinic orbits have the exact parametric representations:

$$\phi(\xi) = \mp \frac{\sqrt{\psi_M\psi_1} \tanh(\omega_1\xi)}{((\psi_M - \psi_1) + \psi_1 \tanh^2(\omega_1\xi))^{\frac{1}{2}}}, \tag{3.9}$$

where $\omega_1 = \frac{1}{2}\sqrt{(\psi_M - \psi_1)\psi_1}$. The two homoclinic orbits have the exact parametric representations:

$$\phi(\xi) = \mp \left(\psi_1 + \frac{2(\psi_M - \psi_1)\psi_1}{\psi_M \cosh(2\omega_1\xi) + 2\psi_1 - \psi_M} \right)^{\frac{1}{2}}. \tag{3.10}$$

(iv) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (h_1, \infty)$, there exists a family of periodic orbits of system (1.8), enclosing six equilibrium points (see Figure 1(d)). These periodic orbits have the same exact parametric representations as (3.4).

3.4. Exact periodic solutions and homoclinic, heteroclinic solutions of system (1.8) when $(\kappa, \omega) \in (L_2)$ in Figure 1(a).

In this case, we have $\phi_1 = \frac{1}{3}\sqrt{6c}, \phi_2 = \sqrt{2c}, h_2 = 0$ and $h_1 = \frac{c^3}{27}$.

(i) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, h_1)$, there exist three families of periodic orbits of system (1.8), enclosing the origin $O(0, 0), E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively (see Figure 1(e)). These periodic orbits have the same exact parametric representations as (3.6) and (3.7).

(ii) Corresponding to the level curves defined by $H(\phi, y) = \frac{c^3}{27}$, there exist two homoclinic orbits enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively, and two heteroclinic orbits enclosing the origin $O(0, 0)$ (see Figure 1(d)). In this case, (3.2) becomes that $\frac{1}{2}\xi = \int_0^\psi \frac{d\psi}{(\frac{2}{3}c - \psi)\sqrt{(\frac{8}{3}c - \psi)\psi}}$ and $\frac{1}{2}\xi = \int^\psi \frac{d\psi}{(\psi - \frac{2}{3}c)\sqrt{(\frac{8}{3}c - \psi)\psi}}$. Therefore, the two heteroclinic orbits have the exact parametric representations:

$$\phi(\xi) = \mp \frac{4\sqrt{c} \tanh(\omega_2\xi)}{(18 + 6 \tanh^2(\omega_2\xi))^{\frac{1}{2}}}, \tag{3.11}$$

where $\omega_2 = \frac{c}{\sqrt{3}}$. The two homoclinic orbits have the exact parametric representations:

$$\phi(\xi) = \mp \left(\frac{2}{3}c + \frac{2c}{2 \cosh(2\omega_2\xi) - 1} \right)^{\frac{1}{2}}. \tag{3.12}$$

(iii) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (h_1, \infty)$, there exists a family of periodic orbits of system (1.8), enclosing six equilibrium points (see Figure 1(e)). These periodic orbits have the same exact parametric representations as (3.4).

3.5. Exact periodic solutions and homoclinic, heteroclinic solutions of system (1.8) when $(\kappa, \omega) \in (III)$ in Figure 1(a).

In this case, we have $h_2 < 0 < h_1$.

(i) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (h_2, 0]$, there exist two families of periodic orbits of system (1.8), enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively (see Figure 1(f)). (3.2) can be written as $\frac{1}{2}\xi = \int_{\psi_b}^{\psi} \frac{d\psi}{\sqrt{(\psi_a - \psi)(\psi - \psi_b)\psi(\psi + \psi_d)}}$. It gives rise to the following exact parametric representations:

$$\phi(\xi) = \mp \frac{\sqrt{\psi_b}}{(1 - \hat{\alpha}_5^2 \text{sn}^2(\Omega_5 \xi, k))^{\frac{1}{2}}}, \tag{3.13}$$

where $\hat{\alpha}_5^2 = 1 - \frac{\psi_b}{\psi_a}, k^2 = \frac{\hat{\alpha}_5^2 \psi_d}{\psi_b + \psi_d}$ and $\Omega_5 = \frac{1}{4} \sqrt{\psi_a(\psi_b + \psi_d)}$.

Especially, the level curves defined by $H(\phi, y) = 0$ are two periodic orbits enclosing the singular points $E_1(-\phi_2, 0)$ and $E_2(\phi_2, 0)$ respectively. In this case, in (3.13), we have $\psi_a = 2(c + 2\sqrt{c\kappa - \omega}), \psi_b = 2(c - 2\sqrt{c\kappa - \omega}), \psi_d = 0, \hat{\alpha}_5^2 = \frac{4\sqrt{c\kappa - \omega}}{c + 2\sqrt{c\kappa - \omega}}, k^2 = 0, \Omega_5 = \frac{1}{2} \sqrt{c^2 - 4c\kappa + 4\omega}$. Hence, the two periodic orbits have the following exact parametric representations:

$$\phi(\xi) = \mp \frac{\sqrt{\psi_b}}{(1 - \hat{\alpha}_5^2 \sin^2(\Omega_5 \xi))^{\frac{1}{2}}}, \tag{3.14}$$

(ii) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, h_1)$, there exist three families of periodic orbits of system (1.8), enclosing the origin $O(0, 0)$ and $E_1(-\phi_2, 0), E_4(\phi_2, 0)$ respectively (see Figure 1(f)). These periodic orbits have the same exact parametric representations as (3.6) and (3.7).

(iii) Corresponding to the level curves defined by $H(\phi, y) = h_1$, there exist two homoclinic orbits enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively, and two heteroclinic orbits enclosing the origin $O(0, 0)$ (see Figure 1(f)). These orbits have the same exact parametric representations as (3.9) and (3.10).

(iv) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (h_1, \infty)$, there exists a family of periodic orbits of system (1.8), enclosing six equilibrium points (see Figure 1(f)). These periodic orbits have the same exact parametric representations as (3.4).

3.6. Exact periodic solutions and homoclinic, heteroclinic solutions of system (1.8) when $(\kappa, \omega) \in (L_3)$ in Figure 1(a).

In this case, we have $\phi_1 = 0, \phi_2 = \frac{2}{3} \sqrt{6c}, h_1 = 0$ and $h_2 = -\frac{8c^3}{27}$.

(i) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (h_2, 0)$, there exist two families of periodic orbits of system (1.8), enclosing $E_1(-\phi_2, 0)$

and $E_4(\phi_2, 0)$ respectively (see Figure 1(g)). These periodic orbits have the same exact parametric representations as (3.13).

(ii) Corresponding to the level curves defined by $H(\phi, y) = 0, h \in (h_2, 0)$, there exist two homoclinic orbits of system (1.8), enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively (see Figure 1(g)). In this case, (3.2) becomes that $\frac{1}{4}\xi = \int_{\psi}^{4c} \frac{d\psi}{\psi\sqrt{(4c-\psi)\psi}}$. Thus, we have the following exact parametric representations:

$$\phi(\xi) = \mp \frac{4\sqrt{c}}{\sqrt{4 + c^2\xi^2}}. \quad (3.15)$$

(iii) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, \infty)$, there exists a family of periodic orbits of system (1.8), enclosing three equilibrium points (see Figure 1(g)). These periodic orbits have the same exact parametric representations as (3.4).

3.7. Exact periodic solutions and homoclinic solutions of system (1.8) when $(\kappa, \omega) \in (IV)$ in Figure 1(a).

In this case, we have $h_2 < 0$.

(i) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (h_2, 0)$, there exist two families of periodic orbits of system (1.8), enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively (see Figure 1(h)). These periodic orbits have the same exact parametric representations as (3.13).

(ii) Corresponding to the level curves defined by $H(\phi, y) = 0, h \in (h_2, 0)$, system (1.8) has two homoclinic orbits, enclosing $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ respectively (see Figure 1(h)). In this case, (3.2) becomes that $\frac{1}{2}\xi = \int_{\psi}^{\psi_M} \frac{d\psi}{\psi\sqrt{(\psi_M-\psi)(\psi+\psi_i)}}$, where $\psi_M = 2(c + 2\sqrt{c\kappa - \omega})$, $\psi_i = 2(-c + 2\sqrt{c\kappa - \omega})$. Hence, we obtain the exact parametric representations of two homoclinic orbits:

$$\phi(\xi) = \mp \left(\frac{2\psi_M\psi_i}{(\psi_M + \psi_i)\cosh(\omega_0\xi) - (\psi_M - \psi_i)} \right)^{\frac{1}{2}} = \mp \left(\frac{8(c\kappa - \omega - \frac{1}{4}c^2)}{2\sqrt{c\kappa - \omega}\cosh(\omega_0\xi) - c} \right)^{\frac{1}{2}}, \quad (3.16)$$

where $\omega_0 = \frac{1}{2}\sqrt{\psi_M\psi_i} = 2\sqrt{c\kappa - \omega - \frac{1}{4}c^2}$.

(iii) Corresponding to the level curves defined by $H(\phi, y) = h, h \in (0, \infty)$, there exists a family of periodic orbits of system (1.8), enclosing three equilibrium points (see Figure 1(g)). These periodic orbits have the same exact parametric representations as (3.4).

4. Exact parametric representations of the solutions of equation (1.3) with the form (1.4)

In this section, we use the exact solutions of system (1.8) derived in Section 3 to give the exact parametric representations of the solutions of equation (1.3) with the form (1.4). Notice that the function $\theta(\xi) = \frac{1}{2}(c - 2\kappa)\xi + \frac{1}{4}\int\phi^2(\xi)d\xi$. Therefore, we first need to calculate the integral $\int\phi^2(\xi)d\xi$, where $\phi(\xi)$ is given by (3.3)-(3.16).

4.1. Exact solutions of equation (1.3) given by the periodic solutions of system (1.8).

(i) Corresponding to the family of the periodic orbits of system (1.8) enclosing the origin $O(0, 0)$ or the global family of periodic orbits enclosing five or three equilibrium points, given by (3.3) or (3.4), we have

$$\begin{aligned} \int \phi^2(\xi)d\xi &= \int \frac{\psi_3 \hat{\alpha}_1^2 \text{sn}^2(\Omega_1 \xi, k) d\xi}{1 - \hat{\alpha}_1^2 \text{sn}^2(\Omega_1 \xi, k)} \\ &= \frac{\psi_3}{\Omega_1} \left[\Pi \left(\arcsin(\text{sn}(\Omega_1 \xi, k)), \hat{\alpha}_1^2, k \right) - F(\arcsin(\text{sn}(\Omega_1 \xi, k)), k) \right], \end{aligned}$$

where F and Π are normal elliptic integrals of the first kind and the third kind respectively, and

$$\begin{aligned} \int \phi^2(\xi)d\xi &= \int \left(\alpha_1 + \frac{\beta_1}{1 + \hat{\alpha}_2 \text{cn}(\Omega_2 \xi, k)} \right) d\xi \\ &= \alpha_1 \xi + \frac{\beta_1}{1 - \hat{\alpha}_2^2} \left[\Pi \left(\arccos(\text{sn}(\Omega_1 \xi, k)), \frac{\hat{\alpha}_2^2}{\hat{\alpha}_2^2 - 1}, k \right) - \hat{\alpha}_2 f_1 \right], \end{aligned}$$

where the function f_1 can be seen in (Byrd and Fridman [3], pp.215).

Similarly, corresponding to the family of the periodic orbits of system (1.8) enclosing the origin $O(0, 0)$ given by (3.6), we have

$$\begin{aligned} \int \phi^2(\xi)d\xi &= \int \frac{\psi_a \hat{\alpha}_3^2 \text{sn}^2(\Omega_3 \xi, k) d\xi}{1 - \hat{\alpha}_3^2 \text{sn}^2(\Omega_3 \xi, k)} \\ &= \frac{\psi_a}{\Omega_3} \left[\Pi \left(\arcsin(\text{sn}(\Omega_3 \xi, k)), \hat{\alpha}_3^2, k \right) - F(\arcsin(\text{sn}(\Omega_3 \xi, k)), k) \right]. \end{aligned}$$

Especially, corresponding to the periodic orbits of system (1.8) enclosing the origin $O(0, 0)$ given by (14_a), we have

$$\begin{aligned} \int \phi^2(\xi)d\xi &= \int \frac{\psi_2 |\hat{\alpha}_3| \sin(\Omega_3 \xi) d\xi}{(1 - \hat{\alpha}_3^2 \sin^2(\Omega_3 \xi))} \\ &= -\psi_2 \xi + 2 \arctan \left(\frac{(2 - \hat{\alpha}_3^2) \tan(\omega_3 \xi) + \hat{\alpha}_3^2}{2\sqrt{1 - \hat{\alpha}_3^2}} \right). \end{aligned}$$

Write that

$$\begin{aligned} \theta_1(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \frac{\psi_3}{4\Omega_1} \left[\Pi \left(\arcsin(\text{sn}(\Omega_1 \xi, k)), \hat{\alpha}_1^2, k \right) - F(\arcsin(\text{sn}(\Omega_1 \xi, k)), k) \right]; \\ \theta_2(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \frac{1}{4}\alpha_1 \xi + \frac{\beta_1}{4(1 - \hat{\alpha}_2^2)} \left[\Pi \left(\arccos(\text{sn}(\Omega_1 \xi, k)), \frac{\hat{\alpha}_2^2}{\hat{\alpha}_2^2 - 1}, k \right) - \hat{\alpha}_2 f_1 \right]; \\ \theta_3(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \frac{\psi_a}{4\Omega_3} \left[\Pi \left(\arcsin(\text{sn}(\Omega_3 \xi, k)), \hat{\alpha}_3^2, k \right) - F(\arcsin(\text{sn}(\Omega_3 \xi, k)), k) \right]; \\ \theta_4(\xi) &= \frac{1}{2}(c - 2\kappa)\xi - \frac{1}{4}\psi_2 \xi + \frac{1}{2} \arctan \left(\frac{(2 - \hat{\alpha}_3^2) \tan(\omega_3 \xi) + \hat{\alpha}_3^2}{2\sqrt{1 - \hat{\alpha}_3^2}} \right). \end{aligned}$$

Thus, we obtain the following exact solutions of equation (1.3):

$$q(x, t) = q_1(x, t) = \frac{\sqrt{\psi_3} \hat{\alpha}_1 \text{sn}(\Omega_1 \xi, k)}{(1 - \hat{\alpha}_1^2 \text{sn}^2(\Omega_1 \xi, k))^{\frac{1}{2}}} \exp(i[\kappa x - \omega t + \theta_1(\xi)]); \quad (4.1)$$

$$q(x, t) = q_2(x, t) = \left(\alpha_1 + \frac{\beta_1}{1 + \hat{\alpha}_2 \operatorname{cn}(\Omega_2 \xi, k)} \right)^{\frac{1}{2}} \exp(i[\kappa x - \omega t + \theta_2(\xi)]); \quad (4.2)$$

$$q(x, t) = q_3(x, t) = \frac{\sqrt{\psi_a} |\hat{\alpha}_3| \operatorname{sn}(\Omega_3 \xi, k)}{(1 - \hat{\alpha}_3^2 \operatorname{sn}^2(\Omega_3 \xi, k))^{\frac{1}{2}}} \exp(i[\kappa x - \omega t + \theta_3(\xi)]); \quad (4.3)$$

$$q(x, t) = q_4(x, t) = \frac{\phi_2 |\hat{\alpha}_3| \operatorname{sn}(\Omega_3 \xi)}{(1 - \hat{\alpha}_3^2 \sin^2(\Omega_3 \xi))^{\frac{1}{2}}} \exp(i[\kappa x - \omega t + \theta_4(\xi)]). \quad (4.4)$$

(ii) Corresponding to the family of the periodic orbits of system (1.8) enclosing the equilibrium points $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ given by (3.7), (3.13) and (3.14), we have

$$\begin{aligned} \int \phi^2(\xi) d\xi &= \int \left(\psi_c + \frac{\psi_b - \psi_c}{1 - \hat{\alpha}_4^2 \operatorname{sn}^2(\Omega_3 \xi, k)} \right) d\xi \\ &= \psi_c \xi + (\psi_b - \psi_c) \Pi(\operatorname{arcsin}(\operatorname{sn}(\Omega_3 \xi, k)), \hat{\alpha}_4^2, k); \end{aligned}$$

$$\int \phi^2(\xi) d\xi = \int \left(\frac{\psi_b}{1 - \hat{\alpha}_5^2 \operatorname{sn}^2(\Omega_5 \xi, k)} \right) d\xi = \psi_b \Pi(\operatorname{arcsin}(\operatorname{sn}(\Omega_5 \xi, k)), \hat{\alpha}_5^2, k);$$

and

$$\begin{aligned} \int \phi^2(\xi) d\xi &= \int \left(\frac{\psi_b}{1 - \hat{\alpha}_5^2 \sin^2(\Omega_5 \xi, k)} \right) d\xi \\ &= \frac{\psi_b}{\Omega_5 \sqrt{1 - \hat{\alpha}_5^2}} \arctan \left(\frac{\sqrt{2 - \hat{\alpha}_5^2} \tan(\Omega_5 \xi) + \hat{\alpha}_5^2}{2\sqrt{1 - \hat{\alpha}_5^2}} \right). \end{aligned}$$

Write that

$$\theta_5(\xi) = \frac{1}{2}(c - 2\kappa + 2\psi_c)\xi + \frac{1}{4}(\psi_b - \psi_c) \Pi(\operatorname{arcsin}(\operatorname{sn}(\Omega_3 \xi, k)), \hat{\alpha}_4^2, k)$$

and

$$\begin{aligned} \theta_6(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \frac{1}{4}\psi_b \Pi(\operatorname{arcsin}(\operatorname{sn}(\Omega_5 \xi, k)), \hat{\alpha}_5^2, k), \\ \theta_7(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \frac{\psi_b}{4\Omega_5 \sqrt{1 - \hat{\alpha}_5^2}} \arctan \left(\frac{\sqrt{2 - \hat{\alpha}_5^2} \tan(\Omega_5 \xi) + \hat{\alpha}_5^2}{2\sqrt{1 - \hat{\alpha}_5^2}} \right). \end{aligned}$$

Therefore, we obtain the following exact solutions of equation (1.3):

$$q(x, t) = q_5(x, t) = \left(\psi_c + \frac{\psi_b - \psi_c}{1 - \hat{\alpha}_4^2 \operatorname{sn}^2(\Omega_3 \xi, k)} \right)^{\frac{1}{2}} \exp(i[\kappa x - \omega t + \theta_5(\xi)]); \quad (4.5)$$

$$q(x, t) = q_6(x, t) = \frac{\sqrt{\psi_b}}{(1 - \hat{\alpha}_5^2 \operatorname{sn}^2(\Omega_5 \xi, k))^{\frac{1}{2}}} \exp(i[\kappa x - \omega t + \theta_6(\xi)]); \quad (4.6)$$

$$q(x, t) = q_7(x, t) = \left(\frac{\psi_b}{1 - \hat{\alpha}_5^2 \sin^2(\Omega_5 \xi, k)} \right)^{\frac{1}{2}} \exp(i[\kappa x - \omega t + \theta_7(\xi)]). \quad (4.7)$$

(iii) Corresponding to the two heteroclinic orbits of system (1.8) enclosing the equilibrium points $O(0, 0)$ given by (3.5), (3.9) and (3.11), we have

$$\int \phi^2(\xi) d\xi = \frac{16c^3}{3} \int \frac{\xi^2}{9 + c^2 \xi^2} = 16 \left[\frac{1}{3} c \xi - \arctan \left(\frac{1}{3} c \xi \right) \right];$$

$$\begin{aligned} \int \phi^2(\xi)d\xi &= \int \frac{\psi_M \psi_1 \tanh^2(\omega_1 \xi)d\xi}{(\psi_M - \psi_1) + \psi_1 \tanh^2(\omega_1 \xi)} \\ &= \psi_1 \xi - 2 \arctan \left(\sqrt{\frac{\psi_1}{\psi_M - \psi_1}} \tanh \left(\frac{1}{2} \xi \right) \right) \end{aligned}$$

and specially, for (3.11),


$$\int \phi^2(\xi)d\xi = \int \frac{16c \tanh^2(\omega_2 \xi)d\xi}{18 + 6 \tanh^2(\omega_2 \xi)} = \frac{2}{3} c \xi - 2 \arctan \left(\frac{1}{\sqrt{3}} \tanh \left(\frac{1}{2} \xi \right) \right).$$

Write that

$$\begin{aligned} \theta_8(x, t) &= \left(\frac{11}{6} c - \kappa \right) \xi - 4 \arctan \left(\frac{1}{3} c \xi \right); \\ \theta_9(x, t) &= \frac{1}{2} \left(c - 2\kappa + \frac{1}{2} \psi_1 \right) \xi - \frac{1}{2} \arctan \left(\sqrt{\frac{\psi_1}{\psi_M - \psi_1}} \tanh \left(\frac{1}{2} \xi \right) \right); \\ \theta_{10}(x, t) &= \frac{1}{2} \left(\frac{4}{3} c - 2\kappa \right) \xi - \frac{1}{2} \arctan \left(\frac{1}{\sqrt{3}} \tanh \left(\frac{1}{2} \xi \right) \right). \end{aligned}$$

These give rise to the following exact solutions of equation (1.3):

$$q(x, t) = q_8(x, t) = \frac{4c^{\frac{3}{2}} \xi}{\sqrt{3(9 + c^2 \xi^2)}} \exp(i[\kappa x - \omega t + \theta_8(\xi)]); \tag{4.8}$$



(a) (b)

Figure 2. The graphs of the real part and imaginary part of solution $q_8(x, t)$

$$q(x, t) = q_9(x, t) = \frac{\sqrt{\psi_M \psi_1} \tanh(\omega_1 \xi)}{((\psi_M - \psi_1) + \psi_1 \tanh^2(\omega_1 \xi))^{\frac{1}{2}}} \exp(i[\kappa x - \omega t + \theta_9(\xi)]); \tag{4.9}$$

$$q(x, t) = q_{10}(x, t) = \frac{4\sqrt{c} \tanh(\omega_2 \xi)}{(18 + 6 \tanh^2(\omega_2 \xi))^{\frac{1}{2}}} \exp(i[\kappa x - \omega t + \theta_{10}(\xi)]). \tag{4.10}$$

(iv) Corresponding to the two homoclinic orbits of system (1.8) enclosing the equilibrium points $E_1(-\phi_2, 0)$ and $E_4(\phi_2, 0)$ given by (3.10), (3.12), (3.15) and (3.16), we have

$$\begin{aligned} \int \phi^2(\xi)d\xi &= \int \left(\psi_1 + \frac{2(\psi_M - \psi_1)\psi_1}{\psi_M \cosh(2\omega_1 \xi) + 2\psi_1 - \psi_M} \right) d\xi \\ &= 2 \arctan \left(\sqrt{\frac{\psi_M - \psi_1}{\psi_1}} \tanh \left(\frac{1}{2} \omega_1 \xi \right) \right); \end{aligned}$$




Figure 3. The graphs of the real part and imaginary part of solution $q_{10}(x, t)$

$$\begin{aligned} \int \phi^2(\xi) d\xi &= \int \left(\frac{2}{3}c + \frac{2c}{2 \cosh(2\omega_2\xi) - 1} \right) d\xi = \frac{2}{3}c\xi + 2 \arctan \left(\sqrt{3} \tanh \left(\frac{c}{2\sqrt{3}}\xi \right) \right); \\ \int \phi^2(\xi) d\xi &= 16c \int \frac{d\xi}{4 + c^2\xi^2} = 8 \arctan \left(\frac{1}{2}c\xi \right); \\ \int \phi^2(\xi) d\xi &= \int \left(\frac{8(c\kappa - \omega - \frac{1}{4}c^2)}{2\sqrt{c\kappa - \omega} \cosh(\omega_0\xi) - c} \right) d\xi \\ &= 4 \arctan \left(\left(\frac{2\sqrt{c\kappa - \omega} + c}{2\sqrt{c\kappa - \omega} - c} \right)^{\frac{1}{2}} \tanh \left(\sqrt{c\kappa - \omega - \frac{1}{4}c^2}\xi \right) \right). \end{aligned}$$

Write that

$$\begin{aligned} \theta_{11}(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \frac{1}{2} \arctan \left(\sqrt{\frac{\psi_M - \psi_1}{\psi_1}} \tanh \left(\frac{1}{2}\omega_1\xi \right) \right); \\ \theta_{12}(\xi) &= \frac{1}{2} \left(\frac{4}{3}c - 2\kappa \right) \xi + \frac{1}{2} \arctan \left(\sqrt{3} \tanh \left(\frac{c}{2\sqrt{3}}\xi \right) \right); \\ \theta_{13}(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + 4 \arctan \left(\frac{1}{2}c\xi \right); \\ \theta_{14}(\xi) &= \frac{1}{2}(c - 2\kappa)\xi + \arctan \left(\left(\frac{2\sqrt{c\kappa - \omega} + c}{2\sqrt{c\kappa - \omega} - c} \right)^{\frac{1}{2}} \tanh \left(\sqrt{c\kappa - \omega - \frac{1}{4}c^2}\xi \right) \right). \end{aligned}$$

Hence, we obtain the exact solutions of equation (1.3) as follows:

$$q(x, t) = q_{11}(x, t) = \left(\psi_1 + \frac{2(\psi_M - \psi_1)\psi_1}{\psi_M \cosh(2\omega_1\xi) + 2\psi_1 - \psi_M} \right)^{\frac{1}{2}} \exp(i[\kappa x - \omega t + \theta_8(\xi)]); \quad (4.11)$$

$$q(x, t) = q_{12}(x, t) = \left(\frac{2}{3}c + \frac{2c}{2 \cosh(2\omega_2\xi) - 1} \right)^{\frac{1}{2}} \exp(i[\kappa x - \omega t + \theta_9(\xi)]); \quad (4.12)$$

$$q(x, t) = q_{13}(x, t) = \frac{4\sqrt{c}}{\sqrt{4 + c^2\xi^2}} \exp(i[\kappa x - \omega t + \theta_{10}(\xi)]); \quad (4.13)$$

$$q(x, t) = q_{14}(x, t) = \left(\frac{8(c\kappa - \omega - \frac{1}{4}c^2)}{2\sqrt{c\kappa - \omega} \cosh(\omega_0\xi) - c} \right)^{\frac{1}{2}} \exp(i[\kappa x - \omega t + \theta_{11}(\xi)]). \quad (4.14)$$




Figure 4. The graphs of the real part and imaginary part of solution $q_{12}(x, t)$




Figure 5. The graphs of the real part and imaginary part of solution $q_{13}(x, t)$

References

- [1] K. K. Al-Kalbani, K. S. Al-Ghafri, E. V. Krishnan and A. Biswas, *Solitons and modulation instability of the perturbed Gerdjikov–Ivanov equation with spatio-temporal dispersion*, *Chaos, Solitons & Fractals*, 2021, 153, Article ID 111523, 8 pages.
- [2] H. M. Baskonus, M. Younis, M. Bilal, U. Younas, S.ur Rehman and W. Gao, *Modulation instability analysis and perturbed optical soliton and other solutions to the Gerdjikov-Ivanov equation in nonlinear optics*, *Modern Physics Letters B*, 2020, 34(35), Article ID 2050404, 15 pages.
- [3] P. F. Byrd, M. D. Fridman, *Handbook of elliptic integrals for engineers and scientists*, Springer, Berlin, 1971.
- [4] H. H. Chen, Y. C. Lee and C. S. Liu, *Integrability of nonlinear Hamiltonian systems by inverse scattering method*, *Physica Scripta*, 1979, 20, 490–492.
- [5] H. H. Dai and E. Fan, *Variable separation and algebro-geometric solutions of the Gerdjikov–Ivanov equation*, *Chaos, Solitons & Fractals*, 2004, 22(1), 93–101.
- [6] M. Dong, L. Tian, J. Wei and Y. Wang, *Some localized wave solutions for the coupled Gerdjikov–Ivanov equation*, *Applied Mathematics Letters*, 2021, 122, Article ID 107483, 7 pages.
- [7] E. Fan, *Integrable evolution systems based on Gerdjikov–Ivanov equations, bi-Hamiltonian structure, finite-dimensional integrable systems and N-fold Darboux transformation*, *Journal of Mathematical Physics*, 2000, 41(11), 7769–7782.
- [8] E. Fan, *Darboux transformation and soliton-like solutions for the Gerdjikov-Ivanov equation*, *Journal of Physics A: Mathematical and General*, 2000, 33, 6925–6933.
- [9] E. Fan, *A family of completely integrable multi-Hamiltonian systems explicitly related to some celebrated equations*, *Journal of Mathematical Physics*, 2001, 42(9), 4327, 19 pages.
- [10] E. Fan, *Integrable systems of derivative nonlinear Schrödinger type and their multi-Hamiltonian structure*, *Journal of Physics A: Mathematical and General*, 2001, 34, 513–519.
- [11] E. Fan, *Bi-Hamiltonian structure and Liouville integrability for a Gerdjikov–Ivanov equation hierarchy*, *Chinese Physics Letters*, 2001, 18(1), 3 pages.
- [12] S. Fromm, *Admissible boundary values for the Gerdjikov-Ivanov equation with asymptotically time-periodic boundary data*, *Symmetry, Integrability and Geometry: Methods and Applications*, 2020, 16, 79, 15 pages.
- [13] V. S. Gerdjikov and M. I. Ivanov, *A quadratic pencil of general type and nonlinear evolution equations, II. Hierarchies of Hamiltonian structures*, *Bulgarian Journal of Physics*, 1983, 10(2), 130–143.
- [14] C. A. S. Gömez, A. Jhangeer, H. Rezazadeh, R. A. Talarposhti and A. Bekir, *Closed form solutions of the perturbed Gerdjikov-Ivanov equation with variable coefficients*, *East Asian Journal on Applied Mathematics*, 2021, 11(1), 207–218.
- [15] B. He and Q. Meng, *Bifurcations and new exact travelling wave solutions for the Gerdjikov–Ivanov equation*, *Communications in Nonlinear Science and Numerical Simulation*, 2010, 15(7), 1783–1790.

- [16] H. Indexed, M. Saleh and A. A. Altwaty, *Optical solitons of the extended Gerdjikov–Ivanov equation in DWDM system by extended simplest equation method*, Applied Mathematics & Information Sciences, 2020, 14(5), 901–907.
- [17] D. J. Kaup and A. C. Newell, *An exact solution for a derivative nonlinear Schrödinger equation*, Journal of Mathematical Physics, 1978, 19(4), 798, 4 pages.
- [18] M. M. A. Khater, *Abundant wave solutions of the perturbed Gerdjikov-Ivanov equation in telecommunication industry*, Modern Physics Letters B, 2021, 35(26), Article ID 2150456, 23 pages.
- [19] J. Li, *Singular nonlinear traveling wave equations: bifurcations and exact solutions*, Science Press, Beijing, 2013.
- [20] J. Li, *Geometric properties and exact travelling wave solutions for the generalized Burger-Fisher equation and the Sharma-Tasso-Olver equation*, Journal of Nonlinear Modeling and Analysis, 2019, 1(1), 1–10.
- [21] M. Li, Y. Zhang, R. Ye and Y. Lou, *Exact solutions of the nonlocal Gerdjikov-Ivanov equation*, Communications in Theoretical Physics, 2021, 73, Article ID 105005, 8 pages.
- [22] Y. Li, L. Zhang, B. Hu and R. Wang, *The initial-boundary value for the combined Schrödinger and Gerdjikov–Ivanov equation on the half-line via the Riemann–Hilbert approach*, Theoretical and Mathematical Physics, 2021, 209, 1537–1551.
- [23] Y. Lou, Y. Zhang, R. Ye and M. Li, *Modulation instability, higher-order rogue waves and dynamics of the Gerdjikov–Ivanov equation*, Wave Motion, 2021, 106, Article ID 102795, 10 pages.
- [24] P. Lu, Y. Wang and C. Dai, *Abundant fractional soliton solutions of a space-time fractional perturbed Gerdjikov-Ivanov equation by a fractional mapping method*, Chinese Journal of Physics, 2021, 74, 96–105.
- [25] J. Luo and E. Fan, *[Math Processing Error]-dressing method for the coupled Gerdjikov–Ivanov equation*, Applied Mathematics Letters, 2020, 110, Article ID 106589, 10 pages.
- [26] J. Luo and E. Fan, *Dbar-dressing method for the Gerdjikov–Ivanov equation with nonzero boundary conditions*, Applied Mathematics Letters, 2021, 120, Article ID 107297, 10 pages.
- [27] T. Tsuchida, *Integrable discretizations of derivative nonlinear Schrödinger equations*, Journal of Physics A: Mathematical and General, 2002, 35(36), 7827–7847.
- [28] Vinita and S. S. Ray, *Lie symmetry reductions, power series solutions and conservation laws of the coupled Gerdjikov–Ivanov equation using optimal system of Lie subalgebra*, Zeitschrift für angewandte Mathematik und Physik, 2021, 72, 133, 18 pages.
- [29] X. Xiao and Z. Yin, *Exact single travelling wave solutions to the fractional perturbed Gerdjikov–Ivanov equation in nonlinear optics*, Modern Physics Letters B, 2021, 35(22), Article ID 2150377, 13 pages.
- [30] S. Zhang, T. Xu, M. Li and X. Zhang, *Higher-order algebraic soliton solutions of the Gerdjikov–Ivanov equation. Asymptotic analysis and emergence of rogue*

-
- waves*, Physica D: Nonlinear Phenomena, 2022, 432, Article ID 133128, 14 pages.
- [31] Z. Zhang and E. Fan, *Inverse scattering transform for the Gerdjikov–Ivanov equation with nonzero boundary conditions*, Zeitschrift für angewandte Mathematik und Physik, 2020, 71, 149, 28 pages.
- [32] Z. Zhang and E. Fan, *Inverse scattering transform and multiple high-order pole solutions for the Gerdjikov–Ivanov equation under the zero/nonzero background*, Zeitschrift für angewandte Mathematik und Physik, 2021, 72(4), 153, 25 pages.
- [33] Y. Zhou and J. Zhuang, *Bifurcations and exact solutions of the Raman soliton model in nanoscale optical waveguides with metamaterials*, Journal of Nonlinear Modeling and Analysis, 2021, 3(1), 145–165.