

# Generalized Coupled Fixed Point Results on Complex Partial Metric Space Using Contractive Condition

Mani Gunaseelan<sup>1</sup>, Lakshmi Narayan Mishra<sup>2</sup> and Vishnu Narayan Mishra<sup>3,†</sup>

**Abstract** In this paper, we obtain coupled fixed point results on complex partial metric space under contractive condition. An example to support our result is presented.

**Keywords** Coupled fixed point, Complex partial metric space.

**MSC(2010)** 47H10, 54H25.

## 1. Introduction

In many branches of science, economics, computer science, engineering and the development of nonlinear analysis, the fixed point theory is one of the most important tool. In 1989, Backhtin [4] introduced the concept of b-metric space. In 1993, Czerwik [6] extended the results of b-metric spaces. Altun, Sola and Simsek [1] introduced generalized contractions on partial metric spaces. Rao, Kishore, Tas, Satyanaraya and Prasad [10] introduced common coupled fixed point results in ordered partial metric spaces. Azam, Fisher and Khan [3] introduced new spaces called complex valued metric spaces and established the existence of fixed point theorems under the contraction condition. Dhivya and Marudai [7] introduced new spaces called complex partial metric space and established the existence of common fixed point theorems under the contraction condition of rational expression. Bhaskar and Lakshmikantham [8] introduced the concept of coupled fixed point. Ćirić and Lakshmikantham [5] investigated some more coupled fixed point theorems in partially ordered sets. Aydi [2] introduced coupled fixed point results on partial metric spaces. Gunaseelan and Mishra [9] introduced coupled fixed point theorems on complex partial metric space using different type of contractive conditions. In this paper, we introduced generalized coupled fixed point results on complex partial metric spaces under the contractive condition.

---

<sup>†</sup>the corresponding author.

Email address: [vishnunarayanmishra@gmail.com](mailto:vishnunarayanmishra@gmail.com)(V. N. Mishra)

<sup>1</sup>Department of Mathematics, Sri Sankara Arts and Science College (Autonomous), Affiliated to Madras University, Enathur, Kanchipuram, Tamil Nadu 631 561, India

<sup>2</sup>Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology (VIT) University, Vellore 632 014, Tamil Nadu, India

<sup>3</sup>Department of Mathematics, Indira Gandhi National Tribal University, Laipur, Amarkantak, Anuppur, Madhya Pradesh 484887, India

## 2. Preliminaries

Let  $\mathbb{C}$  be the set of complex numbers and  $z_1, z_2 \in \mathbb{C}$ . Define a partial order  $\preceq$  on  $\mathbb{C}$  as follows:

$z_1 \preceq z_2$  if and only if  $Re(z_1) \leq Re(z_2)$ ,  $Im(z_1) \leq Im(z_2)$ .

Consequently, one can infer that  $z_1 \preceq z_2$  if one of the following conditions is satisfied:

- (i)  $Re(z_1) = Re(z_2), Im(z_1) < Im(z_2)$ ,
- (ii)  $Re(z_1) < Re(z_2), Im(z_1) = Im(z_2)$ ,
- (iii)  $Re(z_1) < Re(z_2), Im(z_1) < Im(z_2)$ ,
- (iv)  $Re(z_1) = Re(z_2), Im(z_1) = Im(z_2)$ .

In particular, we write  $z_1 \succcurlyeq z_2$  if  $z_1 \neq z_2$  and one of (i), (ii) and (iii) is satisfied and we write  $z_1 \prec z_2$  if only (iii) is satisfied. Notice that

- (a) If  $0 \preceq z_1 \succcurlyeq z_2$ , then  $|z_1| < |z_2|$ ,
- (b) If  $z_1 \preceq z_2$  and  $z_2 \prec z_3$  then  $z_1 \prec z_3$ ,
- (c) If  $a, b \in \mathbb{R}$  and  $a \leq b$  then  $az \preceq bz$  for all  $z \in \mathbb{C}$ .

**Definition 2.1.** [7] A complex partial metric on a non-empty set  $Y$  is a function  $\sigma_c : Y \times Y \rightarrow \mathbb{C}^+$  such that for all  $p, r, s \in Y$ :

- (i)  $0 \preceq \sigma_c(p, p) \preceq \sigma_c(p, r)$  (small self-distances)
- (ii)  $\sigma_c(p, r) = \sigma_c(r, p)$  (symmetry)
- (iii)  $\sigma_c(p, p) = \sigma_c(p, r) = \sigma_c(r, r) \Leftrightarrow p = r$  (equality)
- (iv)  $\sigma_c(p, r) \preceq \sigma_c(p, s) + \sigma_c(s, r) - \sigma_c(s, s)$  (triangularity).

A complex partial metric space is a pair  $(Y, \sigma_c)$  such that  $Y$  is a non empty set and  $\sigma_c$  is complex partial metric on  $Y$ .

For the complex partial metric  $\sigma_c$  on  $Y$ , the function  $d_{\sigma_c} : Y \times Y \rightarrow \mathbb{C}^+$  given by  $\sigma_c^t = 2\sigma_c(p, r) - \sigma_c(p, p) - \sigma_c(r, r)$  is a (usual) metric on  $Y$ . Each complex partial metric  $\sigma_c$  on  $Y$  generates a topology  $\tau_{\sigma_c}$  on  $Y$  with the base family of open  $\sigma_c$ -balls  $\{B_{\sigma_c}(p, \epsilon) : p \in Y, \epsilon > 0\}$ , where  $B_{\sigma_c}(p, \epsilon) = \{r \in Y : \sigma_c(p, r) < \sigma_c(p, p) + \epsilon\}$  for all  $p \in Y$  and  $0 < \epsilon \in \mathbb{C}^+$ .

**Definition 2.2.** [7] Let  $(Y, \sigma_c)$  be a complex partial metric space (CPMS). A sequence  $(p_n)$  in a CPMS  $(Y, \sigma_c)$  is convergent to  $p \in Y$ , if for every  $0 \prec \epsilon \in \mathbb{C}^+$  there is  $N \in \mathbb{N}$  such that for all  $n \in \mathbb{N}$  we get  $p_n \in B_{\sigma_c}(p, \epsilon)$

**Definition 2.3.** [7] Let  $(Y, \sigma_c)$  be a complex partial metric space. A sequence  $(p_n)$  in a CPMS  $(Y, \sigma_c)$  is called Cauchy if there is  $a \in \mathbb{C}^+$  such that for every  $\epsilon \prec 0$  there is  $N \in \mathbb{N}$  such that for all  $n, m \geq N$ ;  $|\sigma_c(p_n, p_m) - a| < \epsilon$ .

**Definition 2.4.** [7] Let  $(Y, \sigma_c)$  be a complex partial metric space (CPMS).

- (1) A CPMS  $(Y, \sigma_c)$  is said to be complete if a Cauchy sequence  $(p_n)$  in  $Y$  converges, with respect to  $\tau_{\sigma_c}$ , to a point  $p \in Y$  such that  $\sigma_c(p, p) = \lim_{n, m \rightarrow \infty} \sigma_c(p_n, p_m)$ .
- (2) A mapping  $H : Y \rightarrow Y$  is said to be continuous at  $p_0 \in Y$  if for every  $\epsilon \prec 0$ , there exists  $\delta > 0$  such that  $H(B_{\sigma_c}(p_0, \delta)) \subset B_{\sigma_c}(H(p_0, \epsilon))$ .

**Lemma 2.1.** [7] Let  $(Y, \sigma_c)$  be a complex partial metric space. A sequence  $\{y_n\}$  is Cauchy sequence in the CPMS  $(Y, \sigma_c)$  then  $\{y_n\}$  is Cauchy in a metric space  $(Y, \sigma_c^t)$ .

**Definition 2.5.** Let  $(Y, \sigma_c)$  be a complex partial metric space (CPMS). Then an element  $(p, r) \in Y \times Y$  is said to be a coupled fixed point of the mapping  $F :$